

Poincaré's Observation and the Origin of Tsallis Generalized Canonical Distributions

C. Vignat* and A. Plastino**

**L.I.S. Grenoble, BP 46, 38402 St. Martin d'Hères cedex, France, vignat@lis.inpg.fr and*

***Facultad de Ciencias Exactas, National University La Plata
and IFLP-CONICET, C. C. 727, 1900 La Plata, Argentina*

Abstract

In this paper, we present some geometric properties of the maximum entropy (MaxEnt) Tsallis-distributions under energy constraint. In the case $q > 1$, these distributions are proved to be marginals of uniform distributions on the sphere; in the case $q < 1$, they can be constructed as conditional distributions of a Cauchy law built from the same uniform distribution on the sphere using a gnomonic projection. As such, these distributions reveal the relevance of using Tsallis distributions in the microcanonical setup: an example of such application is given in the case of the ideal gas.

PACS numbers: 05.40.-a, 05.20.Gg

I. INTRODUCTION

Nonextensive thermostatics is a very active field nowadays, based on the concept of Tsallis' information measure (or entropy) [1]. The concomitant maximum (Tsallis) entropy distributions play a role for the description of non-extensive systems akin of that of Gaussian ones for extensive systems. By Tsallis distributions, we mean the following n -variate probability densities:

- if $\frac{n}{n+2} < q < 1$, with $\beta = \frac{1}{2q-n(1-q)}$ and $A_q = \frac{\Gamma(\frac{1}{1-q})(\beta(1-q))^{n/2}}{\Gamma(\frac{1}{1-q}-\frac{n}{2})\pi^{n/2}|C|^{1/2}}$

$$f_X(X) = A_q (1 - \beta(q-1)x^T C^{-1}x)^{\frac{1}{q-1}}$$
- if $q > 1$ or $q < 0$, with $\beta = \frac{1}{2q-n(1-q)}$ and $A_q = \frac{\Gamma(\frac{q}{q-1}+\frac{n}{2})(\beta(q-1))^{n/2}}{\Gamma(\frac{q}{q-1})\pi^{n/2}|C|^{1/2}}$

$$f_X(X) = A_q (1 - \beta(q-1)x^T C^{-1}x)^{\frac{1}{q-1}}_+$$

where x_+ denotes $\max(x, 0)$.

These distributions are solutions of the following maximization problem:

$$\max_f \frac{1}{1-q} \int_{\mathbb{R}^n} f^q \text{ such that } \int_{\mathbb{R}^n} x x^T f(x) dx = C.$$

In this sense, they are the counterparts of the Gaussian distribution with covariance matrix C which is solution of

$$\max_f \left(- \int_{\mathbb{R}^n} f \log f \right) \text{ such that } \int_{\mathbb{R}^n} x x^T f(x) dx = C$$

Gaussian distributions exhibit universal properties that allow to derive important statistical results such as the central limit theorem or the entropy power inequality. Some of these results have a "Tsallis-counterpart" [1]. A geometric approach like the one advanced here is interesting since it sheds new light on these properties.

II. GEOMETRIC APPROACH

A. stochastic properties

In this section, we propose a geometric characterization of Tsallis distributions based on their stochastic properties. Thus we begin by a detailed stochastic study of Tsallis random vectors. We denote as

$$X \sim T_{q,C}$$

a Tsallis vector with nonextensivity parameter q and covariance matrix C .

case $\frac{n}{n+2} < q < 1$

If $X \sim T_{q,C}$, then a stochastic representation of X writes

$$X = \sqrt{m-2} C^{1/2} \frac{N}{\sqrt{a}}$$

where a is a chi-square distributed random variable with $m = \frac{2}{1-q} - n$ degrees of freedom, independent of the Gaussian random vector N with unit covariance matrix. It is understood that m may be non-integer.

duality result

If $X \sim T_{q,C}$ with $\frac{n}{n+2} < q < 1$ then the following random vector

$$Y = \frac{X}{\sqrt{1 + \frac{1}{m-2} X^T C^{-1} X}}$$

verifies

$$Y \sim T_{q^*, C^*}$$

with

$$C^* = C \frac{m-2}{m+n}, \quad \frac{1}{q^* - 1} = \frac{1}{1-q} - \frac{n}{2} - 1$$

case $q > 1$

From the duality result, we deduce a stochastic representation for $Y \sim T_{q^*, C}$ with $q^* > 1$ as follows

$$Y = \sqrt{m+n} C^{1/2} \frac{N}{\sqrt{a + N^T N}} \quad (1)$$

where a is a chi-square random variable with $m = \frac{2q^*}{q^*-1}$ degrees of freedom, independent of the Gaussian random vector N with unit covariance matrix.

B. Geometric representation

The preceding stochastic representations allow to derive easily a geometric construction of Tsallis random vectors as follows.

case $q > 1$

An n -variate random vector U is uniformly distributed on the ellipsoid $\mathcal{E}_{C,n} = \{Z \in \mathbb{R}^n | Z^T C Z = 1\}$ if and only if it writes

$$U = \frac{C^{-1/2} N}{\sqrt{N^T N}}$$

where N is a Gaussian vector with unit covariance matrix. Comparing this stochastic representation with the Tsallis vector's one (1), the following result can be proved:

Theorem 1 *If $Y \sim T_{q^*,C}$ with $q^* > 1$ or $q^* < 0$ then Y is the n -variate marginal vector of an $(m+n)$ -variate random vector U uniformly distributed on the ellipsoid $\mathcal{E}_{C^{-1},m+n}$ with $m = \frac{2q^*}{q^*-1}$.*

We remark that, according to this characterization, the $(m+n-1)$ -variate marginal random vector is the only one distributed according to a Tsallis law with negative non-extensivity index (namely $q^* = -1$); moreover, the $(m+n-2)$ -variate marginal is uniform inside a $(m+n-2)$ -dimensional ellipsoid and corresponds to a non-extensivity index $q^* = +\infty$. All other lower dimension marginals have finite and positive index $q^* > 1$.

case $q < 1$

For the sake of simplicity, we address here the uncorrelated case $C = I_n$. We note that this is not a loss of generality, since

$$X \sim T_{q,C} \Rightarrow C^{-1/2} X \sim T_{q,I}.$$

Theorem 2 *Assume point P is uniformly distributed on the sphere $\mathcal{E}_{I_n,n}$; consider the intersection M of vector OP with the hyperplane $\mathcal{H} = \{Z \in \mathbb{R}^n | Z_n = 1\}$.¹ Then point M follows a $(n-1)$ -variate Cauchy distribution in $\mathcal{H}$:*

$$f_M(y_1, \dots, y_{n-1}) \propto (1 + \sum_{i=1}^{n-1} y_i^2)^{-\frac{n}{2}}.$$

Moreover, any distribution of point M conditioned on variables $y_{k+1}, \dots, y_{n-1}$ follows a k -variate Tsallis distribution with $m = n - k$ degrees of freedom [10]:

$$f_M(y_1, \dots, y_k | y_{k+1}, \dots, y_{n-1}) \propto (1 + \lambda \sum_{i=1}^k y_i^2)^{-\frac{n}{2}}$$

¹ point M is called the gnomonic projection of point P




FIG. 1: the $n = 3$ dimensional sphere, the Cauchy distribution and one of its conditional laws, the Tsallis $q = \frac{1}{3}$ distribution

In Fig. 1 below, the $n = 3$ dimensional sphere is depicted, with the Cauchy distribution of point M : the white curve represents, up to a constant, the one dimensional conditional density $f_M(y_1|y_2)$ which is Tsallis distributed with $m = 2$ degrees of freedom ($q = \frac{1}{3}$).

III. COMMENTS ON SOME PARTICULAR CASES

In this section, we comment on the geometric results obtained.

A. the case $q > 1$ and Poincaré's observation

The characterization of Tsallis random vectors - with $q > 1$ and identity covariance matrix - as marginals of vectors uniformly distributed on a sphere can be related to a result attributed to Poincaré called "Poincaré's Observation" [2] (for a discussion on this paternity, see [8]).

Theorem 3 *If $X = (x_1, \dots, x_p)$ is uniformly distributed on the sphere S_p with radius $\sqrt{p}$, then*

for fixed $n < +\infty$,

$$\lim_{p \rightarrow \infty} \Pr(\cap_{i=1}^n (a_i \leq x_i \leq b_i)) = \int_{a_1}^{b_1} \frac{e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx \dots \int_{a_n}^{b_n} \frac{e^{-\frac{x^2}{2}}}{\sqrt{2\pi}} dx$$

This result shows that a Gaussian (Tsallis $q = 1$ maximizer) random vector with fixed dimension n can be considered as the marginal vector of an infinite-dimensional vector uniformly distributed on the sphere S_∞ . The results obtained above demonstrate that, in contrast, a Tsallis $q > 1$ random vector of dimension n can be viewed as the marginal of a finite-dimensional vector uniformly distributed on the sphere S_p with

$$p = \frac{2q}{q-1} + n.$$

As $q \rightarrow 1^+$, $p \rightarrow +\infty$ and we recover Poincaré's observation.

B. the case $m = 1, q < 1$

The case $m = 1$ and $q < 1$ corresponds to Cauchy-Lorentz distributions, which are not *stricto sensu* Tsallis distributions since they have no finite covariance matrix. However, they appear in many relevant physical situations [4]. They are defined as

$$f_X(X) = \frac{\Gamma\left(\frac{n+1}{2}\right)}{\pi^{\frac{n+1}{2}}} (1 + X^T X)^{-\frac{n+1}{2}}$$

and have for stochastic representation

$$X = \frac{N}{M},$$

where N is an n -variate Gaussian vector and M is a scalar Gaussian variable, independent of N . Applying the above results, we deduce that they can be obtained as gnomonic projections of uniformly distributed points on the n -sphere, as illustrated in the following figure in the case $n = 3$: as point P describes uniformly the surface of S_3 , point M_2 in the plane $X_3 = 1$ is distributed with Tsallis law $T_{1/3, I_2}$.

IV. APPLICATION: THE IDEAL GAS

We adopt here notations of Minlos [5]. Denote as q_i the velocity of the i -th particle among N of an ideal gas in volume Λ . Let us define the set

$$\Omega_{\Lambda, N, E} = \{(q_1, \dots, q_N) \mid H(q_1, \dots, q_N) = E\}$$




FIG. 2: point M_2 distributed with Tsallis law $T_{1/3, I_2}$ while point M_1 is distributed with Tsallis law T_{-1, I_2}

where the Hamiltonian writes

$$H(q_1, \dots, q_N) = \sum_{i=1}^N \frac{1}{2} m q_i^2.$$

In the microcanonical setup, one assumes that the velocities have a distribution ν uniform on surface $\Omega_{\Lambda, N, E}$:

$$\nu(q_1, \dots, q_N) = \frac{2|\Lambda|^N \pi^{3N/2}}{m \Gamma\left(\frac{3N}{2}\right)} \left(\frac{2E}{N}\right)^{\frac{3N}{2}-1}.$$

Let us now consider a subsystem of the gas, consisting of $N_0 < N$ particles in a volume $\Lambda_0 \subset \Lambda$: their distribution can be easily computed as

$$\nu(q_1, \dots, q_{N_0}) \propto \left(E - \sum_{i=1}^{N_0} \frac{1}{2} m q_i^2\right)_+^{\frac{3}{2}(N-N_0)-1}. \quad (2)$$

A usual approach consists then in taking the thermodynamic limit: assuming the total volume Λ converges to $\mathbb{R}^3$, the number of particle per volume $\frac{N}{|\Lambda|}$ converges to density ρ and the energy per volume $\frac{E}{|\Lambda|}$ converges to energy density e , we deduce

$$\nu(q_1, \dots, q_{N_0}) \propto \exp\left(-\frac{3\rho}{2e} \sum_{i=1}^{N_0} \frac{1}{2} m q_i^2\right)$$

and recover the celebrated Boltzmann distribution of velocities.

If, however, we do not take the thermodynamic limit but remain under the finite-dimensional assumption, then we remark that distribution (2) is an N_0 -variate Tsallis distribution with nonextensivity index

$$q = \frac{N - N_0}{N - N_0 - \frac{2}{3}} > 1. \quad (3)$$

In other terms, the distribution of the finite dimensional ideal gas in the microcanonical ensemble maximizes Tsallis entropy with parameter q as defined by (3). As $N - N_0$ is exactly the dimension of the heat bath seen by the N_0 particles, we may conclude that, in this example, Tsallis entropy appears as the finite-dimensional counterpart of Shannon entropy. This idea was already derived by one of the authors [6] years ago, but without the details we provide here. Moreover, Adib et al. [7] have characterized Tsallis $q > 1$ distributions as marginals of uniform distributions, in the case of more general Hamiltonians verifying an homogeneity property, but they assume the (stronger) ergodic hypothesis to derive this result.

-
- [1] M. Gell-Mann and C. Tsallis, Eds. *Nonextensive Entropy: Interdisciplinary applications* (Oxford University Press, Oxford, 2004); A.R. Plastino and A. Plastino, *Phys. Lett. A* **177** (1993) 177.
 - [2] H. Poincaré, *Calcul des Probabilités*, Gauthier-Villars, 1912
 - [3] H.P. McKean, *Geometry of Differential Space*, The Annals of Probability, 1973, vol.1, n.2, pp.197-206
 - [4] S. Abe, A.K. Rajagopal, Information theoretic approach to statistical properties of multivariate Cauchy-Lorentz distributions, *J. Phys. A: Math. Gen.* **34** (2001)
 - [5] R.A. Minlos, *Introduction to mathematical statistical physics*, Providence, R.I. : American Mathematical Society, (2000)
 - [6] A.R. Plastino, A. Plastino, *Phys. Lett. A*, **193** (1994), 251
 - [7] A.B. Adib, A.A. Moreira, J.S. Andrade Jr., M.P. Almeida, Tsallis thermostatics for finite systems: a Hamiltonian approach, *Physica A* **322** (2003) 276- 284
 - [8] P. Diaconis, D. Freedman, A dozen de Finetti-style results in search of a theory, *Ann. Inst. H. Poincaré Probab. Statist.* **23** n.2, suppl. 397–423 (1987)
 - [9] D. Song, A. K. Gupta, Lp-norm uniform distribution, *proceedings of the american mathematical society*, Vol. 125, Number 2, February 1997, Pages 595-601
 - [10] M. E. Johnson, *Multivariate Statistical Simulation*, Wiley and Sons, (1987)